arXiv:1608.06790v1 [math.CV] 24 Aug 2016

Segal-Bargmann-Fock modules of monogenic functions

Dixan Peña Peña
Dipartimento di Matematica
Politecnico di Milano
Via E. Bonardi 9
20133 Milano, Italy
e-mail: dixanpena@gmail.com

Irene Sabadini
Dipartimento di Matematica
Politecnico di Milano
Via E. Bonardi 9
20133 Milano, Italy
e-mail: irene.sabadini@polimi.it

Franciscus Sommen
Clifford Research Group
Department of Mathematical Analysis
Faculty of Engineering and Architecture
Ghent University
Galglaan 2, 9000 Gent, Belgium
e-mail: fs@cage.ugent.be

Abstract

In this paper we introduce the classical Segal-Bargmann transform starting from the basis of Hermite polynomials and extend it to Clifford algebra-valued functions. Then we apply the results to monogenic functions and prove that the Segal-Bargmann kernel corresponds to the kernel of the Fourier-Borel transform for monogenic functionals. This kernel is also the reproducing kernel for the monogenic Bargmann module.

Keywords: Segal-Bargmann-Fock spaces; Segal-Bargmann transform; monogenic functions; Dirac operators.

Mathematics Subject Classification: 30G35, 42B10, 44A15.

1 Introduction

The Segal-Bargmann transform can be seen as a unitary map from spaces of square-integrable functions to spaces of square-integrable holomorphic functions (see [1], [11], [12]). The latter class of functions is known in the literature under different names: Bargmann or Segal-Bargmann or Fock spaces either in one or several complex variables (see [9], [15]). In this paper we do not enter historical discussion on the nomenclature and we will call them Segal-Bargmann-Fock spaces.

It is natural to consider another higher dimensional extension, namely the one based on monogenic functions with values in a Clifford algebra. Recently, in [7], [8], the authors have pointed out that this approach can be useful in studying quantum systems with internal, discrete degrees of freedom corresponding to nonzero spins.

Square-integrable holomorphic functions are well-known in the literature and they may belong to Hardy or Bergman spaces, whose reproducing kernels are rational functions. In the case of the Segal-Bargmann-Fock spaces the kernel is an exponential function. This reproducing kernel is not bounded and this fact makes some computations in this framework more complicated.

On the other hand, the fact that Segal-Bargmann-Fock spaces are defined on the whole space (either $\mathbb{C}$ or $\mathbb{C}^m$) makes other techniques from Fourier analysis available. From the physics point of view, the (normalized) reproducing kernels of the Segal-Bargmann-Fock spaces are the so-called coherent states in quantum mechanics.

In this paper we consider the higher dimensional framework based on monogenic functions. We can introduce a notion of Segal-Bargmann module (over the Clifford algebra) and of Segal-Bargmann-Fock transform. The fact that we have a Fischer decomposition allows to prove a relation between the projection of the transform onto its monogenic part and the Fourier-Borel kernel. Equipping the Segal-Bargmann-Fock module with the Fischer inner product we can also show that the Segal-Bargmann transform is still an isometry on harmonic polynomials.

The plan of the paper is the following. In Section 2 we revise the classical Segal-Bargmann transform. Then in Section 3 we apply the results of Section 2 to the case of monogenic functions. In Section 4 we study monogenic Bargmann modules. Finally in Section 5 we make summarizing remarks.

2 The classical Segal-Bargmann transform

The Segal-Bargmann transform is a very well-known operator acting from the space $L^2(\mathbb{R})$ and the Fock space which is unitary and allows to identify these two spaces (see [1], [11], [12] and the recent monographs [9], [15]). For any $f \in L^2(\mathbb{R})$, we define the Segal-Bargmann transform of f (sometimes called Segal-Bergmann-Fock transform) as

$$\mathcal{B}[f](z) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \exp\left(-\frac{z^2}{2} + xz - \frac{x^2}{4}\right) f(x)\, dx.$$

Let us consider in $L^2(\mathbb{R})$ the basis $\{\psi_k(x)\}$, where

$$\psi_k(x) = H_k(x)e^{-x^2/4}, \qquad H_k(x) = (-1)^k e^{x^2/2}\partial_x^k e^{-x^2/2}.$$

Note that $H_k(x)$ are the well-known Hermite polynomials. This basis is orthogonal, in fact

$$\langle \psi_\ell, \psi_k \rangle = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \overline{\psi_\ell(x)} \psi_k(x)\, dx = \delta_{\ell k} k!,$$

where $\delta_{\ell k}$ is the Kronecker delta.

Using integration by parts we obtain

$$\begin{aligned}
\mathcal{B}[\psi_k](z) &= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \exp\left(-\frac{(z-x)^2}{2}\right) H_k(x)\, dx \\
&= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \partial_x^k \left(\exp\left(-\frac{z^2}{2} + xz\right)\right) \exp(-x^2/2)\, dx \\
&= \frac{1}{\sqrt{2\pi}} z^k \int_{\mathbb{R}} \exp\left(-\frac{(z-x)^2}{2}\right)\, dx \\
&= z^k.
\end{aligned}$$

Let us consider the so-called Fischer space, namely the space of real analytic functions equipped with the inner product

$$[R, S] = \overline{R(\partial_x)} S(x)|_{x=0}.$$

If we consider instead of the complex variable z a real variable u, we can use the Fischer inner product to get the equality $\left[u^\ell, u^k\right] = \delta_{\ell k} k!$.

The above computations show that the Segal-Bargmann-Fock transform maps $f(x) \mapsto \mathcal{B}[f](u)$ which is an isometry between $L^2(\mathbb{R})$ and the Fischer space.

On the other hand we can consider the space $\mathcal{O}(\mathbb{C})$ of entire holomorphic functions equipped with the inner product

$$\langle f, g\rangle = \frac{1}{\pi}\int_{\mathbb{C}} \exp\left(-|z|^2\right)\overline{f(z)}g(z)\,dxdy, \qquad z = x + iy.$$

We note that the basis $\left\{z^k\right\}$ is orthogonal, in fact:

$$\langle z^\ell, z^k\rangle = \frac{1}{\pi}\int_{\mathbb{C}} \exp\left(-|z|^2\right)\overline{z}^k z^\ell\,dxdy = 0 \quad \text{for } k \neq \ell.$$

Moreover

$$\langle z^k, z^k\rangle = \frac{1}{\pi}\int_0^{2\pi}\int_0^{\infty} \exp\left(-r^2\right) r^{2k}\, rdrd\theta = k!, \quad \text{where } z = re^\theta.$$

Let us define the Segal-Bargmann-Fock space $\mathcal{B}(\mathbb{C})$ as the Hilbert space of entire functions and square-integrable with respect to the Gaussian density, i.e.

$$\frac{1}{\pi}\int_{\mathbb{C}} \exp\left(-|z|^2\right)|f(z)|^2\,dxdy < \infty.$$

Then we have that $\mathcal{B}:\ L^2(\mathbb{R}) \to \mathcal{B}(\mathbb{C})$ is an isometry. Indeed, the Fischer inner product for functions in $\mathbb{R}$ is equal to the Bargmann inner product of the holomorphic extensions.

This discussion can be extended from one to several complex variables. Let us denote by $\underline{x}$ the m-tuple $(x_1, \ldots, x_m) \in \mathbb{R}^m$. We consider the space $L^2(\mathbb{R}^m)$ with the basis $\{\psi_{k_1\ldots k_m}(\underline{x})\}$ where

$$\psi_{k_1\ldots k_m}(\underline{x}) = H_{k_1\ldots k_m}(\underline{x})e^{-|\underline{x}|^2/4}$$

and $H_{k_1\ldots k_m}$ are the Hermite polynomials in $\mathbb{R}^m$ given by

$$H_{k_1\ldots k_m}(\underline{x})e^{-|\underline{x}|^2/2} = (-1)^{k_1+\cdots+k_m}\partial_{x_1}^{k_1}\ldots\partial_{x_m}^{k_m}e^{-|\underline{x}|^2/2}.$$

Definition 2.1. *The Segal-Bargmann-Fock transform of $f \in L^2(\mathbb{R}^m)$ is defined by*

$$\mathcal{B}[f](\underline{z}) = \frac{1}{(2\pi)^{m/2}}\int_{\mathbb{R}^m} \exp\left(-\frac{\underline{z}\cdot\underline{z}}{2} + \underline{x}\cdot\underline{z} - \frac{\underline{x}\cdot\underline{x}}{4}\right) f(\underline{x})\,d\underline{x}, \quad \underline{x}\cdot\underline{z} = \sum_{j=1}^m x_j z_j.$$

Similarly to the one variable case we have that

$$\mathcal{B}[\psi_{k_1\ldots k_m}](\underline{z}) = z_1^{k_1}\ldots z_m^{k_m}.$$

In several variables the Segal-Bargmann-Fock space $\mathcal{B}(\mathbb{C}^m)$ is the Hilbert space of entire functions in $\mathbb{C}^m$ which are square-integrable with respect to the $2m$-dimensional Gaussian density, i.e.

$$\frac{1}{\pi^m}\int_{\mathbb{C}^m} \exp\left(-|\underline{z}|^2\right)|f(\underline{z})|^2\,d\underline{x}d\underline{y} < \infty, \qquad \underline{z} = \underline{x} + i\underline{y}$$

and equipped with the inner product

$$\langle f, g\rangle = \frac{1}{\pi^m}\int_{\mathbb{C}^m} \exp\left(-|\underline{z}|^2\right)\overline{f(\underline{z})}g(\underline{z})\,d\underline{x}d\underline{y}.$$

Finally note that $\|\psi_{k_1\ldots k_m}\|^2 = k_1!\ldots k_m!$. It follows from this that $\mathcal{B}:\ L^2(\mathbb{R}^m) \to \mathcal{B}(\mathbb{C}^m)$ is an isometry.

3 The monogenic case

Let us denote by $\mathbb{R}_m$ the real Clifford algebra generated by m imaginary units $e_1, \ldots, e_m$. The multiplication in this associative algebra is determined by the relations

$$e_j e_k + e_k e_j = -2\delta_{ij}.$$

An element $a \in \mathbb{R}_m$ can be written as

$$a = \sum_A a_A e_A, \quad x_A \in \mathbb{R},$$

where the basis elements $e_A = e_{j_1} \ldots e_{j_k}$ are defined for every subset $A = \{j_1, \ldots, j_k\}$ of $\{1, \ldots, m\}$ with $j_1 < \cdots < j_k$. For the empty set, one puts $e_\emptyset = 1$, the latter being the identity element.

Observe that the dimension of $\mathbb{R}_m$ as a real linear space is 2^m. Furthermore, conjugation in $\mathbb{R}_m$ is given by $\overline{a} = \sum_A a_A \overline{e}_A$, where $\overline{e}_A = \overline{e}_{j_k} \ldots \overline{e}_{j_1}$ with $\overline{e}_j = -e_j$, $j = 1, \ldots, m$.

In the Clifford algebra $\mathbb{R}_m$, we can identify the so-called 1-vectors, namely the linear combinations with real coefficients of the elements e_j, $j = 1, \ldots, m$, with the vectors in the Euclidean space $\mathbb{R}^m$. The correspondence is given by the map $(x_1, \ldots, x_m) \mapsto \underline{x} = x_1 e_1 + \ldots + x_m e_m$ and it is obviously one-to-one.

The norm of a 1-vector $\underline{x}$ is defined as $|\underline{x}|^2 = \sqrt{x_1^2 + \ldots + x_m^2}$ and clearly $\underline{x}^2 = -|\underline{x}|^2$. The product of two 1-vectors $\underline{x} = \sum_{j=1}^m x_j e_j$ and $\underline{y} = \sum_{j=1}^m y_j e_j$ splits into a scalar part and a 2-vector or so-called bivector part:

$$\underline{x}\,\underline{y} = -\langle \underline{x}, \underline{y} \rangle + \underline{x} \wedge \underline{y},$$

where

$$\langle \underline{x}, \underline{y} \rangle = \underline{x} \cdot \underline{y} = -\frac{1}{2}(\underline{x}\,\underline{y} + \underline{y}\,\underline{x}) = \sum_{j=1}^m x_j y_j$$

$$\underline{x} \wedge \underline{y} = \frac{1}{2}(\underline{x}\,\underline{y} - \underline{y}\,\underline{x}) = \sum_{j=1}^m \sum_{k=j+1}^m (x_j y_k - x_k y_j) e_j e_k.$$

The complex Clifford algebra $\mathbb{C}_m$ can be seen as the complexification of the real Clifford algebra $\mathbb{R}_m$, i.e. $\mathbb{C}_m = \mathbb{R}_m \oplus i\,\mathbb{R}_m$. Any complex Clifford number $c \in \mathbb{C}_m$ may be written as $c = a + ib$, $a, b \in \mathbb{R}_m$, leading to the definition of the Hermitian conjugation: $c^\dagger = \overline{a} - i\overline{b}$.

The first order differential operator

$$\partial_{\underline{x}} = \sum_{j=1}^m e_j \partial_{x_j}$$

is called the Dirac operator in $\mathbb{R}^m$. Functions in the kernel of this operator are known as monogenic functions (see e.g. [2], [3], [4], [6]).

Definition 3.1. *A function $f : \Omega \subset \mathbb{R}^m \to \mathbb{C}_m$ defined and continuously differentiable in the open set Ω is said to be (left) monogenic if $\partial_{\underline{x}} f(\underline{x}) = 0$ in Ω. We denote by $\mathcal{M}(\mathbb{R}^m)$ the right $\mathbb{C}_m$-module of monogenic functions in $\mathbb{R}^m$.*

A basic result in Clifford analysis is the so-called Fischer decomposition. Every homogeneous polynomial R_k of degree k can be uniquely decomposed as

$$R_k(\underline{x}) = M_k(\underline{x}) + \underline{x}R_{k-1}(\underline{x}),$$

where M_k, R_{k-1} are homogeneous polynomials and $M_k \in \mathcal{M}(\mathbb{R}^m)$. The monogenic polynomial M_k is called the monogenic part of R_k denoted by $\mathcal{M}(R_k)$.

If f is real analytic function near the origin, then it admits a decomposition of the form $f(\underline{x}) = \sum_{k=0}^{\infty} R_k(\underline{x})$ in an open ball centred at the origin. The monogenic part of f is thus defined by

$$\mathcal{M}\big(f(\underline{x})\big) = \sum_{k=0}^{\infty} \mathcal{M}\big(R_k(\underline{x})\big).$$

Let us recall the definition of the Clifford-Hermite polynomials $H_{s,k}(\underline{x})$. They are polynomials with real coefficients in $\underline{x}$ of degree s satisfying

$$H_{s,k}(\underline{x})e^{-|\underline{x}|^2/2}P_k(\underline{x}) = (-1)^s \partial_{\underline{x}}^s \big(P_k(\underline{x})e^{-|\underline{x}|^2/2}\big),$$

where $P_k(\underline{x})$ denotes a homogeneous polynomial of degree k in $\mathcal{M}(\mathbb{R}^m)$.

Put $\psi_{s,k}(\underline{x}) = H_{s,k}(\underline{x})e^{-|\underline{x}|^2/4}$. The set of functions $\big\{\psi_{s,k}(\underline{x})P_k(\underline{x}) : s,k \in \mathbb{N}\big\}$ is an orthogonal basis for $L^2(\mathbb{R}^m)$ (see [13]).

Theorem 3.2. *Assume that $P_k(\underline{x})$ is a homogeneous polynomial of degree k in $\mathcal{M}(\mathbb{R}^m)$. Then the following formula holds:*

$$\mathcal{B}\big[\psi_{s,k}(\underline{x})P_k(\underline{x})\big](\underline{z}) = \underline{z}^s \mathcal{B}\big[P_k(\underline{x})e^{-|\underline{x}|^2/4}\big](\underline{z}).$$

Proof. The formula follows using integration by parts in higher dimensions:

$$\begin{aligned}
\mathcal{B}\big[\psi_{s,k}(\underline{x})P_k(\underline{x})\big](\underline{z}) &= \frac{1}{(2\pi)^{m/2}} \int_{\mathbb{R}^m} \exp\left(-\frac{\underline{z}\cdot\underline{z}}{2} + \underline{x}\cdot\underline{z} - \frac{\underline{x}\cdot\underline{x}}{4}\right) \psi_{s,k}(\underline{x})P_k(\underline{x})\, d\underline{x} \\
&= \frac{(-1)^s}{(2\pi)^{m/2}} \int_{\mathbb{R}^m} \exp\left(-\frac{\underline{z}\cdot\underline{z}}{2} + \underline{x}\cdot\underline{z}\right) \partial_{\underline{x}}^s \big(P_k(\underline{x})e^{-|\underline{x}|^2/2}\big)\, d\underline{x} \\
&= \frac{1}{(2\pi)^{m/2}} \int_{\mathbb{R}^m} \partial_{\underline{x}}^s \left(\exp\left(-\frac{\underline{z}\cdot\underline{z}}{2} + \underline{x}\cdot\underline{z}\right)\right) P_k(\underline{x})e^{-|\underline{x}|^2/2}\, d\underline{x} \\
&= \underline{z}^s \mathcal{B}\big[P_k(\underline{x})e^{-|\underline{x}|^2/4}\big](\underline{z}).
\end{aligned}$$

□

The next problem is to compute $\mathcal{B}\big[P_k(\underline{x})e^{-|\underline{x}|^2/4}\big]$. To this end we consider an example from which the general result will follow.

Lemma 3.3. *Let $P_k(\underline{x}) = (x_1 - e_1e_2x_2)^k$. Then*

$$\mathcal{B}\big[P_k(\underline{x})e^{-|\underline{x}|^2/4}\big](\underline{z}) = (z_1 - e_1e_2z_2)^k.$$

Proof. Let $Z = x_1 - e_1e_2x_2$ and $2\partial_{\bar{Z}} = \partial_{x_1} - e_1e_2\partial_{x_2}$. Note that

$$e^{-|\underline{x}|^2/2} = \exp\left(-\frac{1}{2}Z\bar{Z}\right) \exp\left(\sum_{j=3}^{m} -\frac{1}{2}x_j^2\right).$$

Thus we have

$$Z^k e^{-|x|^2/2} = \left((-2\partial_{\bar{Z}})^k \exp\left(-\frac{1}{2} Z\bar{Z}\right)\right) \exp\left(\sum_{j=3}^{m} -\frac{1}{2}x_j^2\right)$$
$$= \big(-(\partial_{x_1} - e_1e_2\partial_{x_2})\big)^k e^{-|\underline{x}|^2/2}.$$

Therefore

$$\mathcal{B}\big[P_k(\underline{x})e^{-|\underline{x}|^2/4}\big](\underline{z}) = \frac{1}{(2\pi)^{m/2}} \int_{\mathbb{R}^m} \left((\partial_{x_1} - e_1e_2\partial_{x_2})^k \exp\left(-\frac{\underline{z}\cdot\underline{z}}{2} + \underline{x}\cdot\underline{z}\right)\right) e^{-|\underline{x}|^2/2} d\underline{x}$$
$$= \frac{(z_1 - e_1e_2z_2)^k}{(2\pi)^{m/2}} \int_{\mathbb{R}^m} \exp\left(-\frac{(\underline{z}-\underline{x})\cdot(\underline{z}-\underline{x})}{2}\right) d\underline{x}$$
$$= \frac{(z_1 - e_1e_2z_2)^k}{(2\pi)^{m/2}} \int_{\mathbb{R}^m} \exp\left(-\frac{|\underline{x}|^2}{2}\right) d\underline{x}$$
$$= (z_1 - e_1e_2z_2)^k.$$

□

We can now prove the following important result:

Theorem 3.4. *Suppose that $P_k(\underline{x})$ is a homogeneous polynomial of degree k in $\mathcal{M}(\mathbb{R}^m)$. Then the following formula holds:*

$$\mathcal{B}\big[H_{s,k}(\underline{x})e^{-|\underline{x}|^2/4}P_k(\underline{x})\big](\underline{z}) = \underline{z}^s P_k(\underline{z}).$$

Proof. We show that the formula that we have established for $P_k(\underline{x}) = (x_1 - e_1e_2x_2)^k$ in Lemma 3.3 holds for a general $P_k(\underline{x})$. First of all, due to the Spin(m)-invariance, the result also holds for monogenic plane waves functions of the form

$$(\langle \underline{x}, \underline{t}\rangle - \underline{t}\,\underline{s}\langle \underline{x}, \underline{s}\rangle)^k,$$

where $\underline{t}$ and $\underline{s}$ are orthogonal unit vectors. Then we notice that the space of homogeneous monogenic polynomials of degree k is spanned by finitely many monogenic plane waves $(\langle \underline{x}, \underline{t}\rangle - \underline{t}\,\underline{s}\langle \underline{x}, \underline{s}\rangle)^k$ for some choices of the parameters $(\underline{t}, \underline{s})$. This proves the result. □

4 Monogenic Bargmann modules

We will use the following notations.

Definition 4.1. *We denote by $\mathcal{M}^s(\mathbb{R}^m)$ the right $\mathbb{C}_m$-module of s-monogenic functions on $\mathbb{R}^m$, namely the set of smooth functions in the kernel of $\partial_{\underline{x}}^s$. Next, by $\mathcal{M}^s(\mathbb{C}^m)$ we denote the right $\mathbb{C}_m$-module of s-monogenic functions on $\mathbb{C}^m$, namely the set of entire holomorphic functions in the kernel of $\partial_{\underline{z}}^s$, where $\partial_{\underline{z}} = \sum_{j=1}^m e_j\partial_{z_j}$ is the complexified Dirac operator.*

Note that $\mathcal{M}^s(\mathbb{C}^m)$ is spanned by the set of polynomials of the form $\underline{z}^j P_k(\underline{z})$, $j = 0, \ldots, s-1$, where $P_k(\underline{z})$ is complex spherical monogenics of degree $k \in \mathbb{N}$.

Let us now define the monogenic Bargmann module.

Definition 4.2. *The monogenic Bargmann module $\mathcal{MB}^s(\mathbb{C}^m)$ is defined as*

$$\mathcal{MB}^s(\mathbb{C}^m) = \mathcal{M}^s(\mathbb{C}^m) \cap \mathcal{B}(\mathbb{C}^m),$$

and it is is equipped with the inner product defined in $\mathcal{B}(\mathbb{C}^m)$.

Since $H_{s,k}(\underline{x})P_k(\underline{x})$, $j=0,\dots,s-1$, spans $\mathcal{M}^s(\mathbb{R}^m)$, we have the following:

Proposition 4.3. *The map*

$$\mathcal{B}:\ \mathcal{M}^s(\mathbb{R}^m)e^{-|\underline{x}|^2/4}\cap L^2(\mathbb{R}^m)\to \mathcal{MB}^s(\mathbb{C}^m)$$

is an isometry.

Proof. The proof follows from the previous Theorem 3.4. □

From the Fischer decomposition we obtain

$$\frac{\langle \underline{x},\underline{u}\rangle^k}{k!}=\sum_{s=0}^{k}\underline{x}^s Z_{k,s}(\underline{x},\underline{u})\,\underline{u}^s,$$

where $Z_{k,s}(\underline{x},\underline{u})$ are the so-called zonal spherical monogenics (see [14]). The latter are homogeneous polynomials of degree $k-s$ in $\underline{x}$ and $\underline{u}$ and satisfy the two-sided biregular system $\partial_{\underline{x}}Z_{k,s}(\underline{x},\underline{u})=Z_{k,s}(\underline{x},\underline{u})\partial_{\underline{u}}=0$.

One can also check that

$$Z_{k,s}(\underline{x},\underline{u})=\frac{Z_{k-s,0}(\underline{x},\underline{u})}{\beta_{s,k-s}\cdots\beta_{1,k-s}},\quad k\geq s$$

with $\beta_{2s,k}=-2s$, $\beta_{2s+1,k}=-(2s+2k+m)$. Moreover,

$$\begin{aligned}Z_k(\underline{x},\underline{u})&=Z_{k,0}(\underline{x},\underline{u})\\&=\frac{\Gamma\left(\frac{m}{2}-1\right)}{2^{k+1}\Gamma\left(k+\frac{m}{2}\right)}(|\underline{x}||\underline{u}|)^k\left[(k+m-2)C_k^{\frac{m}{2}-1}(t)+(m-2)\frac{\underline{x}\wedge\underline{u}}{|\underline{x}||\underline{u}|}C_{k-1}^{\frac{m}{2}}(t)\right],\end{aligned}$$

where $C_k^{\alpha}(t)$ denotes the classical Gegenbauer polynomial and $t=\dfrac{\langle \underline{x},\underline{u}\rangle}{|\underline{x}||\underline{u}|}$.

Therefore, the Fischer decomposition of $e^{\langle \underline{x},\underline{u}\rangle}$ has the form

$$e^{\langle \underline{x},\underline{u}\rangle}=\sum_{k=0}^{\infty}\frac{\langle \underline{x},\underline{u}\rangle^k}{k!}=\sum_{k=0}^{\infty}\sum_{s=0}^{k}\underline{x}^s Z_{k,s}(\underline{x},\underline{u})\,\underline{u}^s=\sum_{s=0}^{\infty}\underline{x}^s E_s(\underline{x},\underline{u})\,\underline{u}^s,$$

with $E_s(\underline{x},\underline{u})=\sum_{k=s}^{\infty}Z_{k,s}(\underline{x},\underline{u})$.

We note that $E(\underline{x},\underline{u})=E_0(\underline{x},\underline{u})$ is the monogenic part of $e^{\langle \underline{x},\underline{u}\rangle}$, which is known as the Fourier-Borel kernel. This function has been computed in closed form using Bessel functions in [5] (see also [10]).

Let us again denote by $\mathcal{M}$ the projection of a function onto its monogenic part in the complex monogenic Fischer decomposition.

Proposition 4.4. *Let $f(\underline{x})=g(\underline{x})e^{-|\underline{x}|^2/4}\in L^2(\mathbb{R}^m)$. Then we have*

$$\mathcal{M}\big(\mathcal{B}[f](\underline{z})\big)=\frac{1}{(2\pi)^{m/2}}\int_{\mathbb{R}^m}E(\underline{z},\underline{x})g(\underline{x})e^{-|\underline{x}|^2/2}\,d\underline{x}=[E(\underline{z},\underline{x})^{\dagger},g(\underline{x})],$$

where $[\cdot,\cdot]$ denotes the Fischer inner product. In particular when g is monogenic, we obtain $\mathcal{M}\big(\mathcal{B}[f](\underline{z})\big)=g(\underline{z})$.

Proof. From the above remarks we obtain

$$\mathcal{M}\left(\exp\left(-\frac{\underline{z}\cdot\underline{z}}{2}+\underline{x}\cdot\underline{z}-\frac{\underline{x}\cdot\underline{x}}{4}\right)\right)=\mathcal{M}\big(\exp(\underline{x}\cdot\underline{z})\big)e^{-|\underline{x}|^2/4}=E(\underline{z},\underline{x})e^{-|\underline{x}|^2/4}.$$

The statement follows now using

$$\mathcal{M}\big(\mathcal{B}[f](\underline{z})\big)=\frac{1}{(2\pi)^{m/2}}\int_{\mathbb{R}^m}\mathcal{M}\left(\exp\left(-\frac{\underline{z}\cdot\underline{z}}{2}+\underline{x}\cdot\underline{z}-\frac{\underline{x}\cdot\underline{x}}{4}\right)\right)f(\underline{x})\,d\underline{x}.$$

□

5 Real s-monogenic Bargmann modules

Let us recall that for harmonic polynomials $R(\underline{x}), S(\underline{x})$ we have (see also [6])

$$[R(\underline{x}),S(\underline{x})]=R(\partial_{\underline{x}})^\dagger S(\underline{x})|_{\underline{x}=\underline{0}}=\frac{1}{(2\pi)^{m/2}}\int_{\mathbb{R}^m}e^{-|\underline{x}|^2/2}R(\underline{x})^\dagger S(\underline{x})\,d\underline{x}.$$

For $P_k(\underline{x})\in\mathcal{M}(\mathbb{R}^m)$ we have

$$P_k(\underline{u})=[Z_k(\underline{u},\underline{x})^\dagger,P_k(\underline{x})]=[E(\underline{u},\underline{x})^\dagger,P_k(\underline{x})].$$

From this formula one also obtains in another way that for every monogenic $g(\underline{x})$ such that $g(\underline{x})e^{-|\underline{x}|^2/4}\in L^2(\mathbb{R}^m)$ the following formula holds

$$g(\underline{u})=\frac{1}{(2\pi)^{m/2}}\int_{\mathbb{R}^m}E(\underline{u},\underline{x})g(\underline{x})e^{-|\underline{x}|^2/2}\,d\underline{x}.$$

For general polynomials $R(\underline{x}), S(\underline{x})$ this link with the Fischer inner product no longer holds. However, we have that

$$[R(\underline{x}),S(\underline{x})]=\frac{1}{\pi^m}\int_{\mathbb{C}^m}e^{-|\underline{z}|^2}R(\underline{z})^\dagger S(\underline{z})\,d\underline{z}.$$

For instance, take R,S of the form $z_1^{k_1}\dots z_m^{k_m}$. Moreover, $\mathcal{B}$ is an isometry between $L^2(\mathbb{R}^m)$ and the Segal-Bargmann-Fock module equipped with the above inner product. In particular, if $g\in\mathcal{M}^s(\mathbb{R}^m)$ we have that the map

$$\mathcal{B}:\ g(\underline{x})e^{-|\underline{x}|^2/4}\mapsto h(\underline{z})$$

is an isometry. Note also that h is s-monogenic but it is no longer true that $h(\underline{u})=g(\underline{u})$ for $\underline{u}\in\mathbb{R}^m$.

Let us consider $f\in\mathcal{B}(\mathbb{C}^m)$ and let $\mathcal{M}_s$ be the orthogonal projection onto the submodule of s-monogenic functions $\mathcal{MB}^s(\mathbb{C}^m)$. To be more precise, if

$$f(\underline{z})=\sum_{j=0}^{\infty}\underline{z}^j f_j(\underline{z}),\quad \partial_{\underline{z}}f_j(\underline{z})=0,$$

is the Fischer decomposition of f, then $\mathcal{M}_s\big(f(\underline{z})\big)=\sum_{j=0}^{s-1}\underline{z}^j f_j(\underline{z})$.

Define the function

$$B_s(\underline{z},\underline{x})=\mathcal{M}_s\left(\exp\left(-\frac{\underline{z}\cdot\underline{z}}{2}+\underline{x}\cdot\underline{z}\right)\right).$$

Theorem 5.1. *The following formula holds:*

$$B_s(\underline{z},\underline{x}) = \sum_{j=0}^{s-1} \underline{z}^j \sum_{\ell=0}^{[j/2]} \frac{1}{2^\ell \ell!} E_{j-2\ell}(\underline{z},\underline{x})\underline{x}^{j-2\ell}.$$

Proof. We have that

$$\exp\left(-\frac{\underline{z}\cdot\underline{z}}{2}\right) = \sum_{\ell=0}^{\infty} \frac{\underline{z}^{2\ell}}{2^\ell \ell!}$$

and

$$\exp\left(\underline{x}\cdot\underline{z}\right) = \sum_{k=0}^{\infty} \underline{z}^k E_k(\underline{z},\underline{x})\underline{x}^k.$$

Therefore

$$\begin{aligned}\exp\left(-\frac{\underline{z}\cdot\underline{z}}{2}+\underline{x}\cdot\underline{z}\right) &= \sum_{k,\ell=0}^{\infty} \frac{1}{2^\ell \ell!}\underline{z}^{k+2\ell} E_k(\underline{z},\underline{x})\underline{x}^k \\ &= \sum_{j=0}^{\infty} \underline{z}^j \sum_{\ell=0}^{[j/2]} \frac{1}{2^\ell \ell!} E_{j-2\ell}(\underline{z},\underline{x})\underline{x}^{j-2\ell}\end{aligned}$$

from which the desired result easily follows. □

Remark 5.2. Note also the formula

$$\mathcal{M}_s\left(\mathcal{B}\big[g(\underline{x})e^{-|\underline{x}|^2/4}\big](\underline{z})\right) = \frac{1}{(2\pi)^{m/2}}\int_{\mathbb{R}^m} \mathcal{M}_s\left(\exp\left(-\frac{\underline{z}\cdot\underline{z}}{2}+\underline{x}\cdot\underline{z}\right)\right) g(\underline{x})e^{-|\underline{x}|^2/2}\,d\underline{x}.$$

Next, one may introduce the real s-monogenic Bargmann module, which is the set of $g \in \mathcal{M}^s(\mathbb{R}^m)$ such that $g(\underline{x})e^{-|\underline{x}|^2/4}$ is square-integrable. To put it otherwise, we can now define $L^2\big(\mathbb{R}^m, e^{-|\underline{x}|^2/4}\big)$ with the inner product

$$\langle f, g\rangle = \frac{1}{(2\pi)^{m/2}}\int_{\mathbb{R}^m} f(\underline{x})^\dagger g(\underline{x})e^{-|\underline{x}|/2}\,d\underline{x}$$

and we may put

$$\mathcal{MB}^s(\mathbb{R}^m) = \mathcal{M}^s(\mathbb{R}^m) \cap L^2\big(\mathbb{R}^m, e^{-|\underline{x}|^2/2}\big).$$

For any $f \in \mathcal{MB}^s(\mathbb{R}^m)$ we can define its Weierstrass transform $\mathcal{W}:\ \mathcal{MB}^s(\mathbb{R}^m) \to \mathcal{MB}^s(\mathbb{C}^m)$ by

$$\mathcal{W}[f](\underline{z}) = \frac{1}{(2\pi)^{m/2}}\int_{\mathbb{R}^m} \exp\left(-\frac{(\underline{z}-\underline{x})\cdot(\underline{z}-\underline{x})}{2}\right) f(\underline{x})\,d\underline{x} = \mathcal{B}\big[f(\underline{x})e^{-|\underline{x}|^2/4}\big](\underline{z}).$$

The map $\mathcal{W}:\ \mathcal{MB}^s(\mathbb{R}^m) \to \mathcal{MB}^s(\mathbb{C}^m)$ is an isometry, moreover

$$\mathcal{W}[f](\underline{z}) = \frac{1}{(2\pi)^{m/2}}\int_{\mathbb{R}^m} B_s(\underline{z},\underline{x})f(\underline{x})e^{-|\underline{x}|^2/2}\,d\underline{x}.$$

Acknowledgments

D. Peña Peña acknowledges the support of a Postdoctoral Fellowship given by Istituto Nazionale di Alta Matematica (INdAM) and cofunded by Marie Curie actions.

References

[1] V. Bargmann, *On a Hilbert space of analytic functions and an associated integral transform*, Comm. Pure Appl. Math. 14 (1961) 187–214.

[2] F. Brackx, R. Delanghe, F. Sommen, *Clifford Analysis*, Pitman Res. Notes in Math., 76, 1982.

[3] F. Colombo, I. Sabadini, F. Sommen, D. C. Struppa, *Analysis of Dirac Systems and Computational Algebra*, Progress in Mathematical Physics, Vol. 39, Birkhäuser, Boston, 2004.

[4] R. Delanghe, F. Sommen, V. Souček, *Clifford algebra and spinor-valued functions*, Mathematics and its Applications, 53, Kluwer Academic Publishers Group, Dordrecht, 1992.

[5] N. De Schepper, F. Sommen, *Closed form of the Fourier-Borel kernel in the framework of Clifford analysis*, Results Math. 62 (2012), no. 1-2, 181–202.

[6] J. Gilbert, M. Murray, *Clifford Algebras and Dirac Operators in Harmonic Analysis*, Cambridge Studies in Advanced Mathematics 26, Cambridge University Press, 1991.

[7] W. D. Kirwin, J. Mourão, J. P. Nunes, T. Qian, *Extending coherent state transforms to Clifford analysis*, arXiv:1601.01380.

[8] J. Mourão, J. P. Nunes, T. Qian, *Coherent State Transforms and the Weyl Equation in Clifford Analysis*, arXiv:1607.06233.

[9] Y. A. Neretin, *Lectures on Gaussian Integral Operators and Classical Groups*, EMS Series of Lectures in Mathematics, European Mathematical Society, 2011.

[10] I. Sabadini, F. Sommen, *A Fourier–Borel transform for monogenic functionals on the Lie ball*, to appear in Journal of the Mathematical Society of Japan.

[11] I. E. Segal, *Mathematical characterization of the physical vacuum for a linear Bose-Einstein field*, Illinois J. Math. 6 (1962), 500–523.

[12] I. E. Segal, *The complex wave representation of the free Boson field*, Topics in functional analysis (essays dedicated to M. G. Kreĭn on the occasion of his 70th birthday), pp. 321–343, Adv. in Math. Suppl. Stud., 3, Academic Press, New York-London, 1978.

[13] F. Sommen, *Special functions in Clifford analysis and axial symmetry*, J. Math. Anal. Appl. 130 (1988), no. 1, 110–133.

[14] F. Sommen, B. Jancewicz, *Explicit solutions of the inhomogeneous Dirac equation*, J. Anal. Math. 71 (1997), 59–74.

[15] K. Zhu, *Analysis on Fock spaces*, Graduate Texts in Mathematics 263, Springer New York 2012.